



MEDICAL ALERT

ASTHMA REACTION

Life Threatening If Prompt Treatment Is Not Given

STUDENT'S NAME

Name of School

LIST ALLERGIES TRIGGERS HERE


WARNING SIGNS OF AN ASTHMA REACTION

- | | |
|-------------------------------------|------------------------------|
| *Wheezing | *Abnormal Breathing Patterns |
| *Coughing | *Nasal Flaring |
| *Shortness of Breath | *Chest Pain |
| *Breathing that Requires Extra Work | *Anxiety |
| *Rapid Pulse | *Nasal Flaring |
| *Bluish Color | *Sweating |

EMERGENCY TREATMENT (Create for each student.)

1. Administer 2 puffs of student's rescue inhaler.
2. Follow "Asthma First" instructions on back.
3. After several minutes, if condition has not improved, administer 2 more puffs of student's rescue inhaler.
4. Do not leave student alone and do not send student to the office alone!
5. Call Parent(s)/Guardian(s)
6. If the Asthma Attack is not abating and you have a sense of impending doom, call 911.

HOW TO USE THE EPIPEN AUTO-INJECTOR.....THREE STEPS



1. Pull off gray safety cap.
2. PUSH black tip onto thigh.
3. Hold for 10 seconds. Call 911.